

2011 AP[®] CALCULUS BC FREE-RESPONSE QUESTIONS (Form B)

6. Let $f(x) = \ln(1 + x^3)$.

- (a) The Maclaurin series for $\ln(1 + x)$ is $x - \frac{x^2}{2} + \frac{x^3}{3} - \frac{x^4}{4} + \dots + (-1)^{n+1} \cdot \frac{x^n}{n} + \dots$. Use the series to write the first four nonzero terms and the general term of the Maclaurin series for f .
- (b) The radius of convergence of the Maclaurin series for f is 1. Determine the interval of convergence. Show the work that leads to your answer.
- (c) Write the first four nonzero terms of the Maclaurin series for $f'(t^2)$. If $g(x) = \int_0^x f'(t^2) dt$, use the first two nonzero terms of the Maclaurin series for g to approximate $g(1)$.
- (d) The Maclaurin series for g , evaluated at $x = 1$, is a convergent alternating series with individual terms that decrease in absolute value to 0. Show that your approximation in part (c) must differ from $g(1)$ by less than $\frac{1}{5}$.
-

2006 AP[®] CALCULUS BC FREE-RESPONSE QUESTIONS (Form B)

6. The function f is defined by $f(x) = \frac{1}{1 + x^3}$. The Maclaurin series for f is given by

$$1 - x^3 + x^6 - x^9 + \dots + (-1)^n x^{3n} + \dots,$$

which converges to $f(x)$ for $-1 < x < 1$.

- (a) Find the first three nonzero terms and the general term for the Maclaurin series for $f'(x)$.
- (b) Use your results from part (a) to find the sum of the infinite series $-\frac{3}{2^2} + \frac{6}{2^5} - \frac{9}{2^8} + \dots + (-1)^n \frac{3n}{2^{3n-1}} + \dots$.
- (c) Find the first four nonzero terms and the general term for the Maclaurin series representing $\int_0^x f(t) dt$.
- (d) Use the first three nonzero terms of the infinite series found in part (c) to approximate $\int_0^{1/2} f(t) dt$. What are the properties of the terms of the series representing $\int_0^{1/2} f(t) dt$ that guarantee that this approximation is within $\frac{1}{10,000}$ of the exact value of the integral?
-

WRITE ALL WORK IN THE EXAM BOOKLET.

2010 AP[®] CALCULUS BC FREE-RESPONSE QUESTIONS (Form B)

5. Let f and g be the functions defined by $f(x) = \frac{1}{x}$ and $g(x) = \frac{4x}{1 + 4x^2}$, for all $x > 0$.
- (a) Find the absolute maximum value of g on the open interval $(0, \infty)$ if the maximum exists. Find the absolute minimum value of g on the open interval $(0, \infty)$ if the minimum exists. Justify your answers.
- (b) Find the area of the unbounded region in the first quadrant to the right of the vertical line $x = 1$, below the graph of f , and above the graph of g .
-

6. The Maclaurin series for the function f is given by $f(x) = \sum_{n=2}^{\infty} \frac{(-1)^n (2x)^n}{n-1}$ on its interval of convergence.
- (a) Find the interval of convergence for the Maclaurin series of f . Justify your answer.
- (b) Show that $y = f(x)$ is a solution to the differential equation $xy' - y = \frac{4x^2}{1+2x}$ for $|x| < R$, where R is the radius of convergence from part (a).
-

WRITE ALL WORK IN THE EXAM BOOKLET.

END OF EXAM

AP[®] CALCULUS BC
2009 SCORING GUIDELINES (Form B)

Question 6

The function f is defined by the power series

$$f(x) = 1 + (x+1) + (x+1)^2 + \cdots + (x+1)^n + \cdots = \sum_{n=0}^{\infty} (x+1)^n$$

for all real numbers x for which the series converges.

- (a) Find the interval of convergence of the power series for f . Justify your answer.
- (b) The power series above is the Taylor series for f about $x = -1$. Find the sum of the series for f .
- (c) Let g be the function defined by $g(x) = \int_{-1}^x f(t) dt$. Find the value of $g\left(-\frac{1}{2}\right)$, if it exists, or explain why $g\left(-\frac{1}{2}\right)$ cannot be determined.
- (d) Let h be the function defined by $h(x) = f(x^2 - 1)$. Find the first three nonzero terms and the general term of the Taylor series for h about $x = 0$, and find the value of $h\left(\frac{1}{2}\right)$.
-

2008 AP[®] CALCULUS BC FREE-RESPONSE QUESTIONS

x	$h(x)$	$h'(x)$	$h''(x)$	$h'''(x)$	$h^{(4)}(x)$
1	11	30	42	99	18
2	80	128	$\frac{488}{3}$	$\frac{448}{3}$	$\frac{584}{9}$
3	317	$\frac{753}{2}$	$\frac{1383}{4}$	$\frac{3483}{16}$	$\frac{1125}{16}$

3. Let h be a function having derivatives of all orders for $x > 0$. Selected values of h and its first four derivatives are indicated in the table above. The function h and these four derivatives are increasing on the interval $1 \leq x \leq 3$.
- Write the first-degree Taylor polynomial for h about $x = 2$ and use it to approximate $h(1.9)$. Is this approximation greater than or less than $h(1.9)$? Explain your reasoning.
 - Write the third-degree Taylor polynomial for h about $x = 2$ and use it to approximate $h(1.9)$.
 - Use the Lagrange error bound to show that the third-degree Taylor polynomial for h about $x = 2$ approximates $h(1.9)$ with error less than 3×10^{-4} .
-

2008 AP[®] CALCULUS BC FREE-RESPONSE QUESTIONS (Form B)

6. Let f be the function given by $f(x) = \frac{2x}{1+x^2}$.
- Write the first four nonzero terms and the general term of the Taylor series for f about $x = 0$.
 - Does the series found in part (a), when evaluated at $x = 1$, converge to $f(1)$? Explain why or why not.
 - The derivative of $\ln(1+x^2)$ is $\frac{2x}{1+x^2}$. Write the first four nonzero terms of the Taylor series for $\ln(1+x^2)$ about $x = 0$.
 - Use the series found in part (c) to find a rational number A such that $\left| A - \ln\left(\frac{5}{4}\right) \right| < \frac{1}{100}$. Justify your answer.
-

GO ON TO THE NEXT PAGE.

2005 AP[®] CALCULUS BC FREE-RESPONSE QUESTIONS (Form B)

3. The Taylor series about $x = 0$ for a certain function f converges to $f(x)$ for all x in the interval of convergence. The n th derivative of f at $x = 0$ is given by

$$f^{(n)}(0) = \frac{(-1)^{n+1}(n+1)!}{5^n(n-1)^2} \text{ for } n \geq 2.$$

The graph of f has a horizontal tangent line at $x = 0$, and $f(0) = 6$.

- (a) Determine whether f has a relative maximum, a relative minimum, or neither at $x = 0$. Justify your answer.
(b) Write the third-degree Taylor polynomial for f about $x = 0$.
(c) Find the radius of convergence of the Taylor series for f about $x = 0$. Show the work that leads to your answer.
-

2007 AP[®] CALCULUS BC FREE-RESPONSE QUESTIONS (Form B)

6. Let f be the function given by $f(x) = 6e^{-x/3}$ for all x .

- (a) Find the first four nonzero terms and the general term for the Taylor series for f about $x = 0$.
(b) Let g be the function given by $g(x) = \int_0^x f(t) dt$. Find the first four nonzero terms and the general term for the Taylor series for g about $x = 0$.
(c) The function h satisfies $h(x) = kf'(ax)$ for all x , where a and k are constants. The Taylor series for h about $x = 0$ is given by

$$h(x) = 1 + x + \frac{x^2}{2!} + \frac{x^3}{3!} + \cdots + \frac{x^n}{n!} + \cdots.$$

Find the values of a and k .

WRITE ALL WORK IN THE EXAM BOOKLET.

END OF EXAM